# **FULL PRESCRIBING** INFORMATION

**NOITAMRO3NI** 

**FULL PRESCRIBING** 

# HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AZACITIDINE FOR INJECTION safely and effectively. See full prescribing information for AZACITIDINE FOR INJECTION.

AZACITIDINE for injection, for subcutaneous or intravenous use Initial U.S. Approval: 2004

	•	Hep
RECENT MAJOR CHANGES		for to
Dosage and Administration (2.1) 5/2022	•	Rena
Warnings and Precautions (5.1.5.6) 5/2022		its m

- INDICATIONS AND USAGE Azacitidine for injection is a nucleoside metabolic inhibitor indicated for the treatment of:

- Adult patients with the following FAB myelodysplastic syndrome (MDS) subtypes: Refractory anemia (RA) or refractory anemia with ringed sideroblasts (RARS) (if accompanied by neutropenia or thrombocytopenia or requiring transfusions), refractory anemia with excess blasts (RAEB), refractory anemia with excess blasts in transformation (RAEB-T), and chronic myelomonocytic leukemia (CMMoL). (1.1) --- DOSAGE AND ADMINISTRATION --
- Do not substitute azacitidine for injection for oral azacitidine. The indications and dosing regimen for azacitidine for injection differ from that of oral azacitidine (2.1, 5.1).
- MDS: The recommended starting dosage for the first treatment cycle, for all patients regardless of baseline hematology values, is azacitidine for injection 75 mg/m<sup>2</sup> daily for 7 days to be administered by subcutaneous injection or intravenous infusion. See full prescribing information for schedule for subsequent cycles. Premedicate for nausea and vomiting (2.2).

Continue treatment as long as the patient continues to benefit; (2.3).

- Monitor all patients for hematologic response and for renal toxicity; delay or reduce dosage as appropriate (2.3, 2.6, 2.7). - DOSAGE FORMS AND STRENGTHS ---
- Lyophilized powder in 100 mg single-dose vials (3).
- -- CONTRAINDICATIONS Advanced Malignant Hepatic Tumors (4.1).
- **FULL PRESCRIBING INFORMATION: CONTENTS\***
- INDICATIONS AND USAGE 1.1 Myelodysplastic Syndromes (MDS)
- 2 DOSAGE AND ADMINISTRATION Important Administration Information
  - 2.2 First Treatment Cycle for Adults
  - 2.3 Subsequent Treatment Cycles for Adults

  - 2.5 Dosage Adjustment Based on Hematology Laboratory Values
  - Dosage Adjustment Based on Serum Electrolytes and Renal Toxicity
  - 2.7 Use in Geriatric Patients 2.8 Preparation of Azacitidine for Injection
  - 2.9 Instructions for Subcutaneous Administration
  - 2.10 Instructions for Intravenous Administration
- DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS
- Advanced Malignant Hepatic Tumors
- 4.2 Hypersensitivity to Azacitidine or Mannitol
- WARNINGS AND PRECAUTIONS
- Risks of Substitution with Other Azacitidine Products 5.2 Anemia, Neutropenia and Thrombocytopenia
- Hepatotoxicity in Patients with Severe Pre-existing Hepatic Impairment

Azacitidine for injection is indicated for treatment of adult patients with the following

(RA) or refractory anemia with ringed sideroblasts (if accompanied by neutropenia

or thrombocytopenia or requiring transfusions), refractory anemia with excess blasts

(RAEB), refractory anemia with excess blasts in transformation (RAEB-T), and chronic

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for

injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug

Do not substitute azacitidine for injection for oral azacitidine. The indications and dosing

regimen for azacitidine for injection differ from that of oral azacitidine [see Warnings and

The recommended starting dose for the first treatment cycle, for all patients regardless of

baseline hematology laboratory values, is 75 mg/m<sup>2</sup> subcutaneously or intravenously, daily

Obtain complete blood counts, liver chemistries and serum creatinine prior to the first dose.

Repeat cycles every 4 weeks. The dose may be increased to 100 mg/m² if no beneficial

effect is seen after 2 treatment cycles and if no toxicity other than nausea and vomiting

has occurred. It is recommended that patients be treated for a minimum of 4 to 6 cycles.

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for

injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug

Platelets (x109/L)

Less than 25

25 to 50

Greater than 50

of the next cycle, in which case continue the current dose.

For adult patients whose baseline counts are WBC less than 3 x10<sup>9</sup>/L, ANC

Monitor patients for hematologic response and renal toxicities [see Warnings and

2.5 Dosage Adjustment Based on Hematology Laboratory Values

However, complete or partial response may require additional treatment cycles. Treatment

Precautions (5.4)], and delay or reduce dosage if necessary [see Dosage and Administration

For adult patients with baseline (start of treatment) WBC greater than or equal

to 3 x10<sup>9</sup>/L, ANC greater than or equal to 1.5 x10<sup>9</sup>/L, and platelets greater than

or equal to 75 x10<sup>9</sup>/L, adjust the dose as follows, based on nadir counts for any

less than 1.5 x10<sup>9</sup>/L, or platelets less than 75 x10<sup>9</sup>/L, base dose adjustments on

nadir counts and bone marrow biopsy cellularity at the time of the nadir as noted

below, unless there is clear improvement in differentiation (percentage of mature

granulocytes is higher and ANC is higher than at onset of that course) at the time

% Dose in the Next Course

50%

67%

100%

French-American-British (FAB) myelodysplastic syndrome subtypes: refractory anemia

5.4 Renal Toxicity

**FULL PRESCRIBING INFORMATION** 

1.1 Myelodysplastic Syndromes (MDS)

- Tumor Lysis Syndrome 5.6 Embryo-Fetal Toxicity

INDICATIONS AND USAGE

mvelomonocytic leukemia (CMMoL).

Precautions (5.1)]

product is not labeled with that information.

2.1 Important Administration Information

for 7 days. Premedicate patients for nausea and vomiting

may be continued as long as the patient continues to benefit.

Nadir Counts

2.3 Subsequent Treatment Cycles for Adults

product is not labeled with that information.

given cycle:

ANC (x109/L)

Less than 0.5

0.5 to 1.5

Greater than 1.5

2 DOSAGE AND ADMINISTRATION

2.2 First Treatment Cycle for Adults

Risks of Substitution with Other Azacitidine Products: Do not substitute azacitidine for injection for oral azacitidine (2.1, 5.1). Anemia, Neutropenia and Thrombocytopenia: Monitor complete blood counts (CBC)

- WARNINGS AND PRECAUTIONS

- frequently (5.2).
- patotoxicity: Patients with severe preexisting hepatic impairment are at higher risk toxicity (5.3).
- nal Toxicity: Monitor patients with renal impairment for toxicity since azacitidine and
- metabolites are primarily excreted by the kidneys (5.4).
- Tumor Lysis Syndrome: Azacitidine may cause fatal or serious tumor lysis syndrome, including in patients with MDS. Assess baseline risk and monitor and treat as appropriate (5.5).
- Embryo-Fetal Toxicity: Azacitidine can cause fetal harm. Advise female patients and male patients with female partners of reproductive potential of the potential risk to a fetus and to use effective contraception (5.6) - ADVERSE REACTIONS

Most common adverse reactions (>30%) in adult patients with MDS by subcutaneous route are: nausea, anemia, thrombocytopenia, vomiting, pyrexia, leukopenia, diarrhea, injection site erythema, constipation, neutropenia and ecchymosis. Most common adverse reactions by intravenous route also included petechiae, rigors, weakness and hypokalemia (6.1). To report SUSPECTED ADVERSE REACTIONS, contact Apotex Corp. at 1-800-706-5575 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

--- USE IN SPECIFIC POPULATIONS

Hypersensitivity to Azacitidine or Mannitol (4.2).

Lactation: Advise not to breastfeed (8.2).

### See 17 for PATIENT COUNSELING INFORMATION.

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information.

Revised: 07/2023

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience **USE IN SPECIFIC POPULATIONS**
- 8.1 Pregnancy
- 8.2 Lactation
- Females and Males of Reproductive Potential 8.3
- 8.4 Pediatric Use
- 8.5 Geriatric Use 10 OVERDOSAGE
- DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
  - 12.1 Mechanism of Action 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility CLINICAL STUDIES
- 14.1 Myelodysplastic Syndromes (MDS) 14.2
- 15 REFERENCES
- HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

### Bone Marrow Biopsy Cellularity at Time of Nadir WBC or Platelet Nadir % decrease in counts from baseline 15 to 30 Less than 15 % Dose in the Next Course 50 to 75 Greater than 75

days after the start of the preceding course, provided that both the WBC and the platelet counts are greater than 25% above the nadir and rising. If a greater than 25% increase above the nadir is not seen by day 28, reassess counts every 7 days. If a 25% increase is not seen by day 42, reduce the scheduled dose by

If a nadir as defined in the table above has occurred, give the next course 28

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information.

2.6 Dosage Adjustment Based on Serum Electrolytes and Renal Toxicity

If unexplained reductions in serum bicarbonate levels to less than 20 mEg/L occur, reduce the dosage by 50% for the next course. Similarly, if unexplained elevations of BUN or serum creatinine occur, delay the next cycle until values return to normal or baseline and reduce the dose by 50% for the next course [see Warnings and Precautions (5.4)]. 2.7 Use in Geriatric Patients

Azacitidine and its metabolites are known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, select the dose carefully and monitor renal function [see Warnings and Precautions (5.4) and Use in Specific Populations (8.5)].

2.8 Preparation of Azacitidine for Injection

Azacitidine for injection is a hazardous drug. Follow applicable special handling and disposal procedures.

The azacitidine for injection vial is single-dose and does not contain any preservatives. Discard unused portions of each vial properly [see How Supplied/Storage and Handling (16)]. Do not save any unused portions for later administration. 2.9 Instructions for Subcutaneous Administration

Reconstitute azacitidine for injection aseptically with 4 mL Sterile Water for Injection, USP to obtain a concentration of 25 mg/mL. Inject the diluent slowly into the vial. Vigorously shake or roll the vial until a uniform suspension is achieved. The suspension will be cloudy. Do not filter the suspension after reconstitution. Doing so could remove the active substance. Preparation for Immediate Subcutaneous Administration: For doses requiring more than

1 vial, divide the dose equally between the syringes (e.g., dose 150 mg = 6 mL, 2 syringes with 3 mL in each syringe) and inject into two separate sites. Due to retention in the vial and needle, it may not be feasible to withdraw all of the suspension from the vial. The product may be held at room temperature for up to 1 hour, but must be administered within 1 hour after reconstitution.

Preparation for Delayed Subcutaneous Administration: The reconstituted product may be kept in the vial or drawn into a syringe. For doses requiring more than 1 vial, divide the dose equally between the syringes (e.g., dose 150 mg = 6 mL, 2 syringes with 3 mL in each syringe) and inject into two separate sites. Due to retention in the vial and needle, it may not be feasible to withdraw all of the suspension from the vial. The product must be refrigerated immediately. See Table 2 for suspension stability storage timelines based on the temperature of the diluent for delayed subcutaneous administration. After removal from refrigerated conditions, the suspension may be allowed to equilibrate to

room temperature for up to 30 minutes prior to administration.

**Subcutaneous Administration** To provide a homogeneous suspension, the contents of the dosing syringe must be resuspended immediately prior to administration. To re-suspend, vigorously roll the syringe

from an old site and never into areas where the site is tender, bruised, red, or hard.

between the palms until a uniform, cloudy suspension is achieved. Azacitidine for injection suspension is administered subcutaneously. Rotate sites for each injection (thigh, abdomen, or upper arm). New injections should be given at least one inch

Table 2. Suspension Stability: Storage timelines based on the temperature of the diluent for suspension stability storage

### Suspension Stability Storage timelines Storage Temperature/Duration Store at room temperature at 25° C (77° F) for up to 1 Room temperature (25° C / 77° F) hour or refrigerated at 2° to 8° C (36° to 46° F) for up to Sterile Water for Injection, USP 8 hours Cold (2° to 8° C / 36° to 46° F) Store refrigerated at 2° to 8° C (36° to 46° F) for up to Sterile Water for Injection, USP 22 hours.

### 2.10 Instructions for Intravenous Administration

Parenteral drug product should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use the product if there is evidence of particulate matter or discoloration. Adult Patients with MDS

Reconstitute the appropriate number of azacitidine for injection vials to achieve the desired dose. Reconstitute each vial with 10 mL Sterile Water for Injection, USP. Vigorously shake or roll the vial until all solids are dissolved. The resulting solution will contain azacitidine 10 mg/ mL. The solution should be clear.

Withdraw the required amount of azacitidine for injection solution to deliver the desired dose and inject into a 50 to 100 mL infusion bag of either 0.9% Sodium Chloride Injection, USP or Lactated Ringer's Injection, USP. Intravenous Solution Incompatibility

Azacitidine for injection is incompatible with 5% Dextrose Injection, USP solutions, Hespan, or solutions that contain bicarbonate. These solutions have the potential to increase the rate of degradation of azacitidine for injection and should therefore be avoided.

Intravenous Administration Azacitidine for injection solution is administered intravenously. Administer the total dose over a period of 10 to 40 minutes. The administration must be completed within 1 hour of

reconstitution of the azacitidine for injection vial. Solution Stability: azacitidine for injection reconstituted and diluted for intravenous administration may be stored at 25°C (77°F), but administration must be completed within 1 hour of reconstitution

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information

# 3 DOSAGE FORMS AND STRENGTHS

Azacitidine for injection is supplied as lyophilized powder in 100 mg single-dose vials.

### 4 CONTRAINDICATIONS

4.1 Advanced Malignant Hepatic Tumors

Azacitidine for injection is contraindicated in patients with advanced malignant hepatic tumors [see Warnings and Precautions (5.3)].

### 4.2 Hypersensitivity to Azacitidine or Mannitol Azacitidine for injection is contraindicated in patients with a known hypersensitivity to

azacitidine or mannitol.

# 5 WARNINGS AND PRECAUTIONS

### 5.1 Risks of Substitution with Other Azacitidine Products Due to substantial differences in the pharmacokinetic parameters [see Clinical Pharmacology

(12.3)], the recommended dose and schedule for azacitidine for injection are different from those of oral azacitidine products. Treatment of patients using azacitidine for injection at the recommended dosage of oral azacitidine may result in a fatal adverse reaction. Treatment of patients using oral azacitidine at the doses recommended for azacitidine for injection may not be effective. Do not substitute azacitidine for injection for oral azacitidine [see Dosage and Administration

# 5.2 Anemia, Neutropenia and Thrombocytopenia

product is not labeled with that information.

risk and monitor and treat as appropriate.

Azacitidine causes anemia, neutropenia and thrombocytopenia in adult patients with MDS. Monitor complete blood counts frequently for response and/or toxicity, at a minimum, prior to In adult patients with MDS, after administration of the recommended dosage for the first

cycle, adjust dosage for subsequent cycles based on nadir counts and hematologic response [see Dosage and Administration (2.5)]. Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug

5.3 Hepatotoxicity in Patients with Severe Pre-existing Hepatic Impairment Because azacitidine is potentially hepatotoxic in patients with severe pre-existing hepatic impairment, caution is needed in patients with liver disease. Patients with extensive tumor burden due to metastatic disease have been reported to experience progressive hepati coma and death during azacitidine treatment, especially in such patients with baseline albumin < 30 g/L. Azacitidine is contraindicated in patients with advanced malignant hepatic tumors [see Contraindications (4.1)]. Monitor liver chemistries prior to initiation of therapy and with each cycle.

Safety and effectiveness of azacitidine in patients with MDS and hepatic impairment have not been studied as these patients were excluded from the clinical trials. Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for

injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information. 5.4 Renal Toxicity Renal toxicity ranging from elevated serum creatinine to renal failure and death have

been reported in patients treated with intravenous azacitidine in combination with other chemotherapeutic agents for non-MDS conditions. In addition, renal tubular acidosis, defined as a fall in serum bicarbonate to < 20 mEq/L in association with an alkaline urine and hypokalemia (serum potassium < 3 mEq/L) developed in 5 patients with CML treated with azacitidine and etoposide. Monitor serum creatinine and electrolytes prior to initiation of therapy and with each cycle. If unexplained reductions in serum bicarbonate < 20 mEq/ L or elevations of BUN or serum creatinine occur, reduce or hold the dose [see Dosage and Administration (2.6)].

Patients with renal impairment may be at increased risk for renal toxicity. Also, azacitidine and its metabolites are primarily excreted by the kidney. Therefore, monitor these patients closely for toxicity [see Dosage and Administration (2.6, 2.7)]. Patients with MDS and renal impairment were excluded from the clinical studies. Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for

injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information. 5.5 Tumor Lysis Syndrome Azacitidine may cause fatal or serious tumor lysis syndrome, including in patients with MDS.

Tumor lysis syndrome may occur despite concomitant use of allopurinol. Assess baseline

5.6 Embryo-Fetal Toxicity Based on the mechanism of action and findings in animals, azacitidine can cause fetal harm

when administered to a pregnant woman. Azacitidine administered to pregnant rats via a single intraperitoneal (IP) dose approximating 8% of the recommended human daily dose caused fetal death and anomalies.

Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with azacitidine and for 6 months after the last dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with azacitidine and for 3 months after the last dose [see Use in Specific Populations (8.1, 8.3)].

### ADVERSE REACTIONS

The following adverse reactions are described in other labeling sections:

- o Anemia, Neutropenia and Thrombocytopenia [see Warnings and Precautions
- Hepatotoxicity in Patients with Severe Pre-existing Hepatic Impairment [see Warnings and Precautions (5.3)]
- Renal Toxicity [see Warnings and Precautions (5.4)]
- Tumor Lysis Syndrome [see Warnings and Precautions (5.5)]

### 6.1 Clinical Trials Experience

neutropenia

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure to azacitidine in 443 patients with MDS from 4 clinical studies. Study 1 was a supportive-care controlled trial (subcutaneous administration), Studies 2 and 3 were single arm studies (one with subcutaneous administration and one with intravenous administration), and Study 4 was an international randomized trial (subcutaneous administration) [see Clinical Studies (14.1)].

In Studies 1, 2 and 3, a total of 268 patients were exposed to azacitidine, including 116 exposed for 6 cycles (approximately 6 months) or more and 60 exposed for greater than 12 cycles (approximately one year). Azacitidine was studied primarily in supportive-care controlled and uncontrolled trials (n=150 and n=118, respectively). The population in the subcutaneous studies (n=220) was 23 to 92 years old (mean 66.4 years), 68% male, and 94% white, and had MDS or AML. The population in the intravenous study (n=48) was 35 to 81 years old (mean 63.1 years), 65% male, and 100% white. Most patients received average daily doses between 50 and 100 mg/m<sup>2</sup>. In Study 4, a total of 175 patients with higher-risk MDS (primarily RAEB and RAEB-T

subtypes) were exposed to azacitidine. Of these patients, 119 were exposed for 6 or more cycles, and 63 for at least 12 cycles. The mean age of this population was 68.1 years (ranging from 42 to 83 years), 74% were male, and 99% were white. Most patients received daily azacitidine doses of 75 mg/m<sup>2</sup> Most Commonly Occurring Adverse Reactions (Subcutaneous or Intravenous Route) in

Adult Patients with MDS: nausea, anemia, thrombocytopenia, vomiting, pyrexia, leukopenia, diarrhea, injection site erythema, constipation, neutropenia, ecchymosis. The most common adverse reactions by intravenous route also included petechiae, rigors, weakness and hypokalemia. Adverse Reactions Most Frequently (>2%) Resulting in Clinical Intervention (Subcutaneous

or Intravenous Route) in Adult Patients with MDS: Discontinuation: leukopenia, thrombocytopenia, neutropenia. Dose Held: leukopenia, neutropenia, thrombocytopenia, pyrexia, pneumonia, febrile

Dose Reduced: leukopenia, neutropenia, thrombocytopenia Table 3 presents adverse reactions occurring in at least 5% of patients treated with azacitidine (subcutaneous) in Studies 1 and 2. It is important to note that duration of exposure was longer for the azacitidine-treated group than for the observation group: patients received azacitidine for a mean of 11.4 months while mean time in the observation arm was 6.1 months.

Number (%) of Patients

### Table 3. Most Frequently Observed Adverse Reactions (≥ 5% in All Subcutaneous Azacitidine Treated Patients; Studies 1 and 2)

System Organ Class	All Azacitidine <sup>b</sup>	Observation <sup>c</sup>	Pyrexia	53 (30)	18 (18)	8 (5)	1 (1)
Preferred Term <sup>a</sup>	(N=220)	(N=92)	Infections and infestations				
Blood and lymphatic system disorders	, ,	, ,	Rhinitis	10 (6)	1 (1)	0	0
Anemia	153 (70)	59 (64)	Upper respiratory tract	16 (9)	4 (4)	3 (2)	0
Anemia aggravated	12 (6)	5 (5)	infection	10 (9)	4 (4)	3 (2)	U
Febrile neutropenia	36 (16)	4 (4)	Urinary tract infection	15 (9)	3 (3)	3 (2)	0
Leukopenia	106 (48)	27 (29)	Investigations				
Neutropenia	71 (32)	10 (11)	Weight decreased	14 (8)	0	1 (1)	0
Thrombocytopenia	144 (66)	42 (46)	Metabolism and nutrition				
Gastrointestinal disorders	(55)	.= (,	disorders				
Abdominal tenderness	26 (12)	1 (1)	Hypokalemia	11 (6)	3 (3)	3 (2)	3 (3)
Constipation	74 (34)	6 (7)	Nervous system disorders				
Diarrhea	80 (36)	13 (14)	Lethargy	13 (7)	2 (2)	0	1 (1)
Gingival bleeding	21 (10)	4 (4)	Psychiatric disorders				
Loose stools	12 (6)	0	Anxiety	9 (5)	1 (1)	0	0
Mouth hemorrhage	11 (5)	1 (1)	Insomnia	15 (9)	3 (3)	0	0
Nausea	155 (71)	16 (17)	Renal and urinary	, ,			
Stomatitis	17 (8)	0	disorders				
Vomiting	, ,	_	Hematuria	11 (6)	2 (2)	4 (2)	1 (1)
General disorders and administration site condition	119 (54)	5 (5)	Respiratory, thoracic and	, ,		, ,	, ,
		F (F)	mediastinal disorders				
Chest pain	36 (16)	5 (5)	Dyspnea	26 (15)	5 (5)	6 (3)	2 (2)
Injection site bruising	31 (14)	0	Dyspnea exertional	9 (5)	1 (1)	)	Ů,
Injection site erythema	77 (35)	0	Pharyngolaryngeal pain	11 (6)	3 (3)	0	0
Injection site granuloma	11 (5)	0	Skin and subcutaneous	( )	. ,		
Injection site pain	50 (23)	0	tissue disorders				
Injection site pigmentation changes	11 (5)	0	Erythema	13 (7)	3 (3)	0	0
Injection site pruritus	15 (7)	0	Petechiae	20 (11)	4 (4)	2 (1)	0
Injection site reaction	30 (14)	0	Pruritus	21 (12)	2 (2)	0	0
Injection site swelling	11 (5)	0	Rash	18 (10)	1 (1)	0	0
Lethargy	17 (8)	2 (2)	Vascular disorders	10 (10)	. ( . )		
Malaise	24(11)	1 (1)	Hypertension	15 (9)	4 (4)	2 (1)	2 (2)
Pyrexia	114 (52)	28 (30)	Multiple reports of the sam				
Infections and infestations			each treatment.	e preferred ter	iii iioiii a patieiit	. Were only cou	inted once within
Nasopharyngitis	32 (15)	3 (3)	edon a edanient.				
Pneumonia	24 (11)	5 (5)	In Studies 1, 2 and 4 with subcutaneous administration of azacitidine, adverse reactions of neutropenia, thrombocytopenia, anemia, nausea, vomiting, diarrhea, constipation,				
Upper respiratory tract infection	28 (13)	4 (4)					
Injury, poisoning, and procedural complications			and injection site erythema/re	action tended	to increase in in	cidence with hi	igher doses of
Post procedural hemorrhage	13 (6)	1 (1)	azacitidine. Adverse reactions that tended to be more pronounced during the first 1 to 2				
Metabolism and nutrition disorders			cycles of subcutaneous treatr	ment compare	d with later cycle	s included thro	ombocytopenia,
Anorexia	45 (21)	6 (7)	neutropenia, anemia, nausea, vomiting, injection site erythema/pain/bruising/reaction,				
Musculoskeletal and connective tissue disorders		, ,	constipation, petechiae, dizzi				
Arthralgia	49 (22)	3 (3)	to be any adverse reactions t				
Chest wall pain	11 (5)	0	Overall, adverse reactions we				
Myalgia	35 (16)	2 (2)	subcutaneous studies. Adverse reactions that appeared to be specifically associated with the intravenous route of administration included infusion site reactions (e.g. erythema or pain)				
Nervous system disorders	(/	\-/				, , ,	ythema or pain)
Dizziness	41 (19)	5 (5)	and catheter site reactions (e				
Headache	48 (22)	10 (11)	In clinical studies of either sul				
Psychiatric disorders	10 (22)	( )	adverse reactions occurring a	at a rate of < 5	% (and not desci	ribed in Tables	∠ or 3) were
Anxiety	29 (13)	3 (3)	reported:	m diaardara	naronula si tos!-	hono marror 4	failura
Allaloty	20 (10)	J (J)	Blood and lymphatic syster	ıı aisoraers:	agranulocytosis,	none marrow i	iaiiure,

Respiratory, thoracic and mediastinal disorders 11 (12) 64 (29) Dyspnea Skin and subcutaneous tissue disorders Dry skin 67 (31) 14 (15) Ecchymosis 37 (17) 4 (4) Erythema 9 (10) 31 (14) Skin nodule 11 (5) 1 (1) 13 (6) 1 (1) Urticaria Vascular disorders Hematoma 19 (9) 15 (7) Hypotension 2 (2) 52 (24) Petechiae 8 (9)

Multiple terms of the same preferred terms for a patient are only counted once within each treatment group.

Includes adverse reactions from all patients exposed to azacitidine, including patients after crossing over from observations. Includes adverse reactions from observation period only; excludes any adverse events

after crossover to azacitidine.

Table 4 presents adverse reactions occurring in at least 5% of patients treated with

azacitidine in Study 4. Similar to Studies 1 and 2 described above, duration of exposure to treatment with azacitidine was longer (mean 12.2 months) compared with best supportive care (mean 7.5 months).

Number (%) of Patients

Grade 3/4

### Table 4. Most Frequently Observed Adverse Reactions (≥ 5% in the Azacitidine Treated Patients and the Percentage with NCI CTC Grade 3/4 Reactions; Study 4)

Any Grade

	, (11	y Grade	5	ade 3/4
System Organ Class Preferred Term <sup>a</sup>	Azacitidine (N=175)	Best Supportive Care Only (N=102)	Azacitidine (N=175)	Care Only (N=102)
Blood and lymphatic system disorders				
Anemia	90 (51)	45 (44)	24 (14)	9 (9)
Febrile neutropenia	24 (14)	10 (10)	22 (13)	7 (7)
Leukopenia	32 (18)	2 (2)	26 (15)	1 (1)
Neutropenia	115 (66)	29 (28)	107 (61)	22 (22)
Thrombocytopenia	122 (70)	35 (34)	102 (58)	29 (28)
Gastrointestinal disorders	(* - )	()	(1-)	()
Abdominal pain	22 (13)	7 (7)	7 (4)	0
Constipation	88 (50)	8 (8)	2 (1)	0
Dyspepsia	10 (6)	2 (2)	0	0
Nausea	84 (48)	12 (12)	3 (2)	0
Vomiting	47 (27)	7 (7)	0	0
General disorders and administration site conditions	()		-	
Fatigue	42 (24)	12 (12)	6 (3)	2 (2)
Injection site bruising	9 (5)	0	0	0
Injection site erythema	75 (43)	0	0	0
Injection site hematoma	11 (6)	0	0	0
Injection site induration	9 (5)	0	0	0
Injection site pain	33 (19)	0	0	0
Injection site rash	10 (6)	0	0	0
Injection site reaction	51 (29)	0	1 (1)	0
Pyrexia	53 (30)	18 (18)	8 (5)	1 (1)
Infections and infestations				
Rhinitis	10 (6)	1 (1)	0	0
Upper respiratory tract infection	16 (9)	4 (4)	3 (2)	0
Urinary tract infection Investigations	15 (9)	3 (3)	3 (2)	0
Weight decreased	14 (8)	0	1 (1)	0
Metabolism and nutrition disorders				
Hypokalemia	11 (6)	3 (3)	3 (2)	3 (3)
Nervous system disorders				
Lethargy	13 (7)	2 (2)	0	1 (1)
Psychiatric disorders				
Anxiety	9 (5)	1 (1)	0	0
Insomnia	15 (9)	3 (3)	0	0
Renal and urinary disorders				
Hematuria	11 (6)	2 (2)	4 (2)	1 (1)
Respiratory, thoracic and mediastinal disorders				
Dyspnea	26 (15)	5 (5)	6 (3)	2 (2)
Dyspnea exertional	9 (5)	1 (1)	0	0
Pharyngolaryngeal pain	11 (6)	3 (3)	0	0
Skin and subcutaneous tissue disorders				
Erythema	13 (7)	3 (3)	0	0
Petechiae	20 (11)	4 (4)	2 (1)	0
Pruritus	21 (12)	2 (2)	0	0
Rash	18 (10)	1 (1)	0	0
Vascular disorders	. ,	• •		
Vasculai disolacis	4 = (0)	4 (4)	2 (1)	2 (2)
Hypertension	15 (9)	+ (+ <i>)</i>	- ( ' '	

Eye disorders: eye hemorrhage

Gastrointestinal disorders: diverticulitis, gastrointestinal hemorrhage, melena, perirectal

General disorders and administration site conditions: catheter site hemorrhage, general physical health deterioration, systemic inflammatory response syndrome.

Hepatobiliary disorders: cholecystitis. **Immune system disorders:** anaphylactic shock, hypersensitivity.

Infections and infestations: abscess limb, bacterial infection, cellulitis, blastomycosis, injection site infection, Klebsiella sepsis, neutropenic sepsis, pharyngitis streptococcal, pneumonia Klebsiella, sepsis, septic shock, Staphylococcal bacteremia, Staphylococcal infection, toxoplasmosis

**Metabolism and nutrition disorders:** dehydration.

Musculoskeletal and connective tissue disorders: bone pain aggravated, muscle

Neoplasms benign, malignant and unspecified: leukemia cutis.

Nervous system disorders: cerebral hemorrhage, convulsions, intracranial hemorrhage. Renal and urinary disorders: loin pain, renal failure.

Respiratory, thoracic and mediastinal disorders: hemoptysis, lung infiltration, pneumonitis, respiratory distress

Skin and subcutaneous tissue disorders: pyoderma gangrenosum, rash pruritic, skin

Surgical and medical procedures: cholecystectomy.

Vascular disorders: orthostatic hypotension

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug

6.2 Postmarketing Experience

product is not labeled with that information.

The following adverse reactions have been identified during postmarketing use of azacitidine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Interstitial lung disease
- Tumor lysis syndrome Injection site necrosis
- Sweet's syndrome (acute febrile neutrophilic dermatosis)
- Necrotizing fasciitis (including fatal cases)
- Differentiation syndrome Pericardial effusion
- Pericarditis

### 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy Risk Summary

Based on its mechanism of action and findings in animals, azacitidine can cause fetal harm when administered to a pregnant woman [see Clinical Pharmacology (12.1)]. There are no data on the use of azacitidine in pregnant women. Azacitidine was teratogenic and caused embryo-fetal lethality in animals at doses lower than the recommended human daily dose (see Data). Advise pregnant women of the potential risk to the fetus.

The background rate of major birth defects and miscarriage is unknown for the indicated population. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

# Animal Data

Early embryotoxicity studies in mice revealed a 44% frequency of intrauterine embryonal death (increased resorption) after a single IP (intraperitoneal) injection of 6 mg/m<sup>2</sup> (approximately 8% of the recommended human daily dose on a mg/m<sup>2</sup> basis) azacitidine on gestation day 10. Developmental abnormalities in the brain have been detected in mice given azacitidine on or before gestation day 15 at doses of ~3 to 12 mg/m² (approximately 4% to 16% the recommended human daily dose on a mg/m<sup>2</sup> basis).

In rats, azacitidine was clearly embryotoxic when given IP on gestation days 4 to 8 (postimplantation) at a dose of 6 mg/m² (approximately 8% of the recommended human daily dose on a mg/m<sup>2</sup> basis), although treatment in the preimplantation period (on gestation days 1 to 3) had no adverse effect on the embryos. Azacitidine caused multiple fetal abnormalities in rats after a single IP dose of 3 to 12 mg/m² (approximately 8% the recommended human daily dose on a mg/m<sup>2</sup> basis) given on gestation day 9, 10, 11 or 12. In this study azacitidine caused fetal death when administered at 3 to 12 mg/m<sup>2</sup> on gestation days 9 and 10; average live animals per litter was reduced to 9% of control at the highest dose on gestation day 9. Fetal anomalies included: CNS anomalies (exencephaly/encephalocele), limb anomalies (micromelia, club foot, syndactyly, oligodactyly), and others (micrognathia, gastroschisis,

# 8.2 Lactation

edema, and rib abnormalities).

There is no information regarding the presence of azacitidine in human milk, the effects of azacitidine on the breastfed infant, or the effects of azacitidine on milk production. Because many drugs are excreted in human milk and because of the potential for tumorigenicity shown for azacitidine in animal studies [see Nonclinical Toxicology (13.1)] and the potential for serious adverse reactions in nursing infants from azacitidine, advise patients not to

### breastfeed during treatment with azacitidine and for 1 week after the last dose. 8.3 Females and Males of Reproductive Potential

Based on its mechanism of action and findings in animals, azacitidine can cause fetal harm when administered to a pregnant woman [see Use in Specific Populations (8.1)].

Pregnancy Testing Verify the pregnancy status of females of reproductive potential prior to initiating azacitidine. Contraception

# <u>Females</u> Advise pregnant women of the potential risk to a fetus. Advise females of reproductive

potential to use effective contraception during treatment with azacitidine and for 6 months after the last dose.

Advise males with female partners of reproductive potential to use effective contraception during treatment with azacitidine and for 3 months after the last dose. Infertility

Based on animal data, azacitidine could have an effect on male or female fertility [see

Nonclinical Toxicology (13.1)]. 8.4 Pediatric Use Safety and effectiveness of azacitidine in pediatric patients with MDS have not been

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information.

# 8.5 Geriatric Use

established.

Of the total number of patients in Studies 1, 2 and 3, 62% were 65 years and older and 21% were 75 years and older. No overall differences in effectiveness were observed between these patients and younger patients. In addition, there were no relevant differences in the frequency of adverse reactions observed in patients 65 years and older compared to

Of the 179 patients randomized to azacitidine in Study 4, 68% were 65 years and older and 21% were 75 years and older. Survival data for patients 65 years and older were consistent

with overall survival results. The majority of adverse reactions occurred at similar frequencies in patients < 65 years of age and patients 65 years of age and older. Elderly patients are more likely to have decreased renal function. Monitor renal function in


these patients [see Dosage and Administration (2.7) and Warnings and Precautions (5.4)].

### 10 OVERDOSAGE

One case of overdose with azacitidine was reported during clinical trials. A patient experienced diarrhea, nausea, and vomiting after receiving a single intravenous dose of approximately 290 mg/m<sup>2</sup>, almost 4 times the recommended starting dose. The events resolved without sequelae, and the correct dose was resumed the following day. In the event of overdosage, the patient should be monitored with appropriate blood counts and should receive supportive treatment, as necessary. There is no known specific antidote for azacitidine overdosage.

# 11 DESCRIPTION

Azacitidine for injection contains azacitidine, which is a nucleoside metabolic inhibitor. Azacitidine is 4-amino-1-β-D-ribofuranosyl-s-triazin-2(1H)-one. The structural formula is as



The empirical formula is C<sub>8</sub>H<sub>12</sub>N<sub>4</sub>O<sub>5</sub>. The molecular weight is 244. Azacitidine is a white to off-white solid. Azacitidine was found to be insoluble in acetone, ethanol, and methyl ethyl ketone; slightly soluble in ethanol/water (50/50), propylene glycol, and polyethylene glycol; sparingly soluble in water, water saturated octanol, 5% dextrose in water, N-methyl-2-pyrrolidone, normal saline and 5% Tween 80 in water; and soluble in dimethylsulfoxide

The finished product is supplied in a sterile form for reconstitution as a suspension for subcutaneous injection or reconstitution as a solution with further dilution for intravenous infusion. Vials of azacitidine for injection contain 100 mg of azacitidine and 100 mg mannitol as a sterile lyophilized powder.

### 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Azacitidine is a pyrimidine nucleoside analog of cytidine. Azacitidine is believed to exert its antineoplastic effects by causing hypomethylation of DNA and direct cytotoxicity on abnormal hematopoietic cells in the bone marrow. The concentration of azacitidine required for maximum inhibition of DNA methylation in vitro does not cause major suppression of DNA synthesis. Hypomethylation may restore normal function to genes that are critical for differentiation and proliferation. The cytotoxic effects of azacitidine cause the death of rapidly dividing cells, including cancer cells that are no longer responsive to normal growth control mechanisms. Non-proliferating cells are relatively insensitive to azacitidine.

# 12.2 Pharmacodynamics

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information. 12.3 Pharmacokinetics

The pharmacokinetics of azacitidine were studied in 6 adult patients with MDS following a single 75 mg/m<sup>2</sup> subcutaneous dose and a single 75 mg/m<sup>2</sup> intravenous dose. Absorption

Azacitidine is rapidly absorbed after subcutaneous administration; the peak plasma azacitidine concentration of 750 ± 403 ng/ml occurred in 0.5 hour after subcutaneous administration. Distribution

The bioavailability of subcutaneous azacitidine relative to intravenous azacitidine is approximately 89%, based on area under the curve. Mean volume of distribution following intravenous dosing is 76 ± 26 L. Mean apparent subcutaneous clearance is 167 ± 49 L/ hour and mean half-life after subcutaneous administration is 41 ± 8 minutes. The AUC and C<sub>max</sub> of subcutaneous administration of azacitidine in 21 patients with cancer were approximately dose proportional within the 25 to 100 mg/m<sup>2</sup> dose range. Multiple dosing at the recommended dose-regimen does not result in drug accumulation with intravenous or subcutaneous administration.

# Elimination

An in vitro study of azacitidine incubation in human liver fractions indicated that azacitidine is not metabolized by the cytochrome P450 (CYP) enzymes. Azacitidine undergoes spontaneous hydrolysis and deamination mediated by cytidine deaminase.

Published studies indicate that urinary excretion is the primary route of elimination of azacitidine and its metabolites. Following intravenous administration of radioactive azacitidine to 5 cancer patients, the cumulative urinary excretion was 85% of the radioactive dose. Fecal excretion accounted for < 1% of administered radioactivity over 3 days. Mean excretion of radioactivity in urine following subcutaneous administration of <sup>14</sup>C-azacitidine was 50%. The mean elimination half-lives of total radioactivity (azacitidine and its metabolites) were similar after intravenous and subcutaneous administrations, about 4

# Specific Populations

The effects of hepatic impairment, gender, or race/ethnicity on the pharmacokinetics of intravenous and subcutaneous azacitidine have not been studied. Pediatric Patients

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information. Patients with Renal Impairment

In adult patients with cancer, the pharmacokinetics of azacitidine in 6 patients with normal renal function (CLcr > 80 mL/min) and 6 patients with severe renal impairment (CLcr < 30 mL/min) were compared following daily subcutaneous dosing (Days 1 through 5) at 75 mg/ m²/day. Severe renal impairment increased azacitidine exposure by approximately 70% after single and 41% after multiple subcutaneous administrations. This increase in exposure was not correlated with an increase in adverse events. The exposure was similar to exposure in patients with normal renal function receiving 100 mg/m<sup>2</sup>.

### Drug-Drug Interactions No formal clinical drug interaction studies with azacitidine have been conducted.

In vitro Studies Cytochrome P450 (CYP) Enzymes: An in vitro study at azacitidine concentrations up to 100

μM (IV Cmax = 10.6 μM) in human liver microsomes indicated that azacitidine does not cause any inhibition of CYP isoforms CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4, or CYP2E1 at clinically achievable concentrations.

In vitro studies with human cultured hepatocytes indicate that azacitidine at concentrations of 1  $\mu$ M to 100  $\mu$ M does not induce CYP 1A2, 2C19, or 3A4/5.

Transporter Systems: An in vitro study with LLC-PK1 cells expressing P-glycoprotein (P-gp) indicated that azacitidine is not a substrate or inhibitor of P-gp.

Azacitidine does not inhibit, breast cancer resistance protein (BCRP), organic anion transporters (OAT) OAT1 and OAT3, organic anion transporting polypeptides (OATP) OATP1B1 and OATP1B3, or organic cation transporter (OCT) OCT2 at clinically relevant

### 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The potential carcinogenicity of azacitidine was evaluated in mice and rats. Azacitidine induced tumors of the hematopoietic system in female mice at 2.2 mg/kg (6.6 mg/m², approximately 8% the recommended human daily dose on a mg/m<sup>2</sup> basis) administered IP three times per week for 52 weeks. An increased incidence of tumors in the lymphoreticular system, lung, mammary gland, and skin was seen in mice treated with azacitidine IP at 2 mg/kg (6 mg/m<sup>2</sup>, approximately 8% the recommended human daily dose on a mg/m<sup>2</sup> basis) once a week for 50 weeks. A tumorigenicity study in rats dosed twice weekly at 15 or 60 mg/m<sup>2</sup> (approximately 20% to 80% the recommended human daily dose on a mg/m<sup>2</sup> basis) revealed an increased incidence of testicular tumors compared with controls. The mutagenic and clastogenic potential of azacitidine was tested in in vitro bacterial systems Salmonella typhimurium strains TA100 and several strains of trpE8, Escherichia coli strains WP14 Pro, WP3103P, WP3104P, and CC103; in in vitro forward gene mutation assay in mouse lymphoma cells and human lymphoblast cells; and in an in vitro micronucleus assay in mouse L5178Y lymphoma cells and Syrian hamster embryo cells. Azacitidine was mutagenic in bacterial and mammalian cell systems. The clastogenic effect of azacitidine was shown by the induction of micronuclei in L5178Y mouse cells and Syrian hamster embryo cells

Administration of azacitidine to male mice at 9.9 mg/m<sup>2</sup> (approximately 9% the recommended human daily dose on a mg/m² basis) daily for 3 days prior to mating with untreated female mice resulted in decreased fertility and loss of offspring during subsequent embryonic and postnatal development. Treatment of male rats 3 times per week for 11 or 16 weeks at doses of 15 to 30 mg/m<sup>2</sup> (approximately 20% to 40%, the recommended human daily dose on a mg/m² basis) resulted in decreased weight of the testes and epididymides, and decreased sperm counts accompanied by decreased pregnancy rates and increased loss of embryos in mated females. In a related study, male rats treated for 16 weeks at 24 mg/m<sup>2</sup> resulted in an increase in abnormal embryos in mated females when examined on day 2 of gestation.

# 14 CLINICAL STUDIES

14.1 Myelodysplastic Syndromes (MDS) Study 1 was a randomized, open-label, controlled trial carried out in 53 U.S. sites compared the safety and efficacy of subcutaneous azacitidine plus supportive care with supportive care alone ("observation") in adult patients with any of the five FAB subtypes of myelodysplastic syndromes (MDS); refractory anemia (RA), RA with ringed sideroblasts (RARS), RA with excess blasts (RAEB), RAEB in transformation (RAEB-T), and chronic myelomonocytic leukemia (CMMoL). RA and RARS patients were included if they met one or more of the following criteria: required packed RBC transfusions; had platelet counts ≤ 50 x 10<sup>9</sup>/ L; required platelet transfusions; or were neutropenic (ANC < 1 x 10<sup>9</sup>/L) with infections requiring treatment with antibiotics. Patients with acute myelogenous leukemia (AML) were not intended to be included. Supportive care allowed in this study included blood transfusion products, antibiotics, antiemetics, analgesics and antipyretics. The use of hematopoietic growth factors was prohibited. Baseline patient and disease characteristics are summarized in Table 6: the 2 groups were similar.

Azacitidine was administered at a subcutaneous dose of 75 mg/m<sup>2</sup> daily for 7 days every 4 weeks. The dose was increased to 100 mg/m<sup>2</sup> if no beneficial effect was seen after 2 treatment cycles. The dose was decreased and/or delayed based on hematologic response or evidence of renal toxicity. Patients in the observation arm were allowed by protocol to cross over to azacitidine if they had increases in bone marrow blasts, decreases in hemoglobin, increases in red cell transfusion requirements, or decreases in platelets, or if they required a platelet transfusion or developed a clinical infection requiring treatment with antibiotics. For purposes of assessing efficacy, the primary endpoint was response rate (as defined in Table 7).

Of the 191 patients included in the study, independent review (adjudicated diagnosis) found that 19 had the diagnosis of AML at baseline. These patients were excluded from the primary analysis of response rate, although they were included in an intent-to-treat (ITT) analysis of all patients randomized. Approximately 55% of the patients randomized to observation crossed over to receive azacitidine treatment.

Table 6. Baseline Demographics and Disease Characteristics

	Azacitidine (N=99)	Observation (N=92)
Gender (n %)		
Male	72 (72.7)	60 (65.2)
Female	27 (27.3)	32 (34.8)
Race (n%)		
White	93 (93.9)	85 (92.4)
Black	1 (1)	1 (1.1)
Hispanic	3 (3)	5 (5.4)
Asian/Oriental	2 (2)	1 (1.1)
Age (years)		
N	99	91
Mean ± SD	67.3 ±10.39	68 ± 10.23
Range	31 to 92	35 to 88
Adjudicated MDS diagnosis at study entry (n%)		
RA	21 (21.2)	18 (19.6)
RARS	6 (6.1)	5 (5.4)
RAEB	38 (38.4)	39 (42.4)
RAEB-T	16 (16.2)	14 (15.2)
CMMoL	8 (8.1)	7 (7.6)
AML	10 (10.1)	9 (9.8)
Transfusion product used in 3 months before study entry (n%)		
Any transfusion product	70 (70.7)	59 (64.1)
Blood cells, packed human	66 (66.7)	55 (59.8)
Platelets, human blood	15 (15.2)	12 (13)
Hetastarch	0 (0)	1 (1.1)
Plasma protein fraction	1 (1)	0 (0)
Other	2 (2)	2 (2.2)

# Table 7. Response Criteria

		RA	RARS	RAEB	RAEB-T	CMMoL
Complete	Marrow	< 5% blasts				
Response (CR), duration ≥ 4 weeks	Peripheral Blood	Normal CBC if abnormal at baseline Absence of blasts in the peripheral circulation				
	Marrow	No marrow		≥ 50% decrease in blasts		
		requiremen	ts	Improvement of marrow dyspoiesis		
Partial Response		≥ 50% restoration in the deficit from normal levels of baseline white cells, hemoglobin and platelets if abnormal at baseline				
(PR),						
duration ≥ 4	Peripheral					
weeks	Blood	No blasts in the peripheral circulation				
		For CMMoL, if WBC is elevated at baseline, a ≥ 75%				
		reduction in the excess count over the unner limit of norr				

The overall response rate (CR + PR) of 15.7% in azacitidine-treated patients without AML (16.2% for all azacitidine randomized patients including AML) was statistically significantly higher than the response rate of 0% in the observation group (p<0.0001) (Table 8). The majority of patients who achieved either CR or PR had either 2 or 3 cell line abnormalities at baseline (79%; 11/14) and had elevated bone marrow blasts or were transfusion dependent at baseline. Patients responding to azacitidine had a decrease in bone marrow blasts percentage, or an increase in platelets, hemoglobin or WBC. Greater than 90% of the responders initially demonstrated these changes by the 5th treatment cycle. All patients who had been transfusion dependent became transfusion independent during PR or CR. The mean and median duration of clinical response of PR or better was estimated as 512 and 330 days, respectively; 75% of the responding patients were still in PR or better at completion of treatment. Response occurred in all MDS subtypes as well as in patients with adjudicated baseline diagnosis of AML.

### Table 8. Response Rates

-	Azacitidine (N=89)	Observation Before Crossover (N=83)		Storage Store unrecon [See USP Co
esponse	n (%)	n (%)	P value	Handling an
verall (CR+PR)	14 (15.7)	0 (0)	(<0.0001)	Azacitidine fo
omplete (CR)	R) 5 (5.6) 0 (0)		(0.06)	procedures. 1
artial (PR)	9 (10.1)	0 (0)		17 PATIEN
				II IAILI


Patients in the observation group who crossed over to receive azacitidine treatment (47 patients) had a response rate of 12.8%.

Study 2, a multi-center, open-label, single-arm study of 72 patients with RAEB, RAEB-T, CMMoL, or AML was also carried out. Treatment with subcutaneous azacitidine resulted in a response rate (CR + PR) of 13.9%, using criteria similar to those described above. The mean and median duration of clinical response of PR or better was estimated as 810 and 430 days, respectively; 80% of the responding patients were still in PR or better at the time of completion of study involvement. In Study 3, another open-label, single-arm study of 48 patients with RAEB, RAEB-T, or AML, treatment with intravenous azacitidine resulted in a response rate of 18.8%, again using criteria similar to those described above. The mean and median duration of clinical response of PR or better was estimated as 389 and 281 days, respectively; 67% of the responding patients were still in PR or better at the time of completion of treatment. Response occurred in all MDS subtypes as well as in patients with adjudicated baseline diagnosis of AML in both of these studies. Azacitidine dosage regimens in these 2 studies were similar to the regimen used in the controlled study. Benefit was seen in patients who did not meet the criteria for PR or better, but were considered "improved." About 24% of azacitidine-treated patients were considered improved and about 2/3 of those lost transfusion dependence. In the observation group, only 5/83 patients met criteria for improvement; none lost transfusion dependence. In all 3 studies, about 19% of patients met criteria for improvement with a median duration of 195 days. Study 4 was an international, multicenter, open-label, randomized trial in patients with MDS with RAEB, RAEB-T or modified CMMoL according to FAB classification and Intermediate-2 and High risk according to IPSS classification. Of the 358 patients enrolled in the study, 179 were randomized to receive azacitidine plus best supportive care (BSC) and 179 were randomized to receive conventional care regimens (CCR) plus BSC (105 to BSC alone, 49 to low dose cytarabine and 25 to chemotherapy with cytarabine and anthracycline). The primary efficacy endpoint was overall survival.

The azacitidine and CCR groups were comparable for baseline parameters. The median age of patients was 69 years (range was 38 to 88 years), 98% were Caucasian, and 70% were male. At baseline, 95% of the patients were higher risk by FAB classification: RAEB (58%), RAEB-T (34%), and CMMoL (3%). By IPSS classification, 87% were higher risk: Int-2 (41%), High (47%). At baseline, 32% of patients met WHO criteria for AML Azacitidine was administered subcutaneously at a dose of 75 mg/m<sup>2</sup> daily for 7 consecutive days every 28 days (which constituted one cycle of therapy). Patients continued treatment until disease progression, relapse after response, or unacceptable toxicity. Azacitidine

patients were treated for a median of 9 cycles (range 1 to 39), BSC only patients for a median of 7 cycles (range 1 to 26), low dose cytarabine patients for a median of 4.5 cycles (range 1 to 15), and chemotherapy with cytarabine and anthracycline patients for a median of 1 cycle (range 1 to 3, i.e. induction plus 1 or 2 consolidation cycles). In the Intent-to-Treat analysis, patients treated with azacitidine demonstrated a statistically significant difference in overall survival as compared to patients treated with CCR (median survival of 24.5 months vs. 15 months; stratified log-rank p=0.0001). The hazard ratio describing this treatment effect was 0.58 (95% CI: 0.43, 0.77).

Kaplan-Meier Curve of Time to Death from Any Cause: (Intent-to-Treat Population)



Key: AZA = azacitidine; CCR = conventional care regimens; CI = confidence interval; HR = Hazard Ratio

Azacitidine treatment led to a reduced need for red blood cell transfusions (see Table 9). In patients treated with azacitidine who were RBC transfusion dependent at baseline and became transfusion independent, the median duration of RBC transfusion independence was 13 months.

Table 9. Effect of Azacitidine on RBC Transfusions in Patients with MDS

Efficacy Parameter	Azacitidine plus BSC (n=179)	Conventional Care Regimens (n=179)
Number and percent of patients who were transfusion dependent at	50/111 (45%)	13/114 (11.4%)
baseline who became transfusion independent on treatment <sup>1</sup>	(95% CI: 35.6%, 54.8%)	(95% CI: 6.2%, 18.7%)
Number and percent of patients who were transfusion-independent at	10/68 (14.7%)	28/65 (43.1%)
baseline who became transfusion- dependent on treatment	(95% CI: 7.3%, 25.4%)	(95% CI: 30.9%, 56%)

A patient was considered RBC transfusion independent during the treatment period if the patient had no RBC transfusions during any 56 consecutive days or more during the treatment period. Otherwise, the patient was considered transfusion dependent.

Pediatric use information is approved for Celgene Corporation's Vidaza (azacitidine for injection). However, due to Celgene Corporation's marketing exclusivity rights, this drug product is not labeled with that information.

### 15 REFERENCES

1. "OSHA Hazardous Drugs." OSHA. http://www.osha.gov/SLTC/hazardousdrugs/index.html

### 16 HOW SUPPLIED/STORAGE AND HANDLING **How Supplied**

Azacitidine for injection is supplied as a lyophilized powder in 100 mg single-dose vials packaged in cartons of 1 vial (NDC 60505-6271-1).

### Store unreconstituted vials at 25° C (77° F); excursions permitted to 15° to 30° C (59° to 86° F) [See USP Controlled Room Temperature].

**Handling and Disposal** Azacitidine for injection is a hazardous drug. Follow applicable special handling and disposal

### 17 PATIENT COUNSELING INFORMATION

Hepatotoxicity in Patients with Severe Pre-Existing Hepatic Impairment

Instruct patients to inform their physician about any underlying liver disease [see Warnings and Precautions (5.3) Renal Toxicity

Instruct patients to inform their physician about any underlying renal disease [see Warnings and Precautions (5.4)]. Embryo-Fetal Toxicity

Advise pregnant women of the potential risk to a fetus [see Warnings and Precautions (5.6) and Use in Specific Populations (8.1)]. Advise females of reproductive potential to use effective contraception during treatment

with azacitidine for injection and for 6 months after the last dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with azacitidine for injection and for 3 months after the last dose. Advise patients to report known or suspected pregnancy to their physicians immediately [see Warnings and Precautions (5.6) and Use in Specific Populations (8.3)], <u>Lactation</u>

Advise patients to avoid breastfeeding while receiving azacitidine for injection and for 1 week after the last dose [see Use in Specific Populations (8.2)].

Advise males and females that azacitidine for injection may impair fertility [see Use in Specific Populations (8.3) and Nonclinical Toxicology (13.1)].

# Manufactured by:

Distributed by:

Revised: July 2023

Jiangsu Hansoh Pharmaceutical Group Co., Ltd., Lianyungang, Jiangsu 222069, China

Apotex Corp. Weston Florida USA 33326