
Drug Facts

Active ingredient (in each tablet)**Purpose**

Loratadine 10 mg..... Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
-

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 6 years and over	1 tablet daily: not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- safety sealed: do not use if induction seal, with “Lift N Peel” tab, under cap is broken or missing
- store between 2°C and 30°C (36°F and 86°F)
- protect from excessive moisture

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose

Questions or comments?

Call **1-800-706-5575**, weekdays, 8:30 am – 5:00 pm Eastern Standard Time

Revised: July 2018